

# Iron (Monoferric, Ferinject and Venofer) Order Form **\*TWO PAGES\***

Send completed form to Coverdale Infusion Clinics: Fax:1-888-236-3502 or by email: [enrollment@coverdaleclinic.com](mailto:enrollment@coverdaleclinic.com)

An infusion fee will apply and is payable by credit card to the Coverdale Resource Center at the time of booking. Please note that a cancellation fee may also apply.

PATIENT INFORMATION			
Patient Name:		DOB (dd/mm/yyyy):	
Patient Allergies:		Patient Address:	
Patient Phone Number: ( ) -		Patient Email:	
Emergency Contact Name:		Patient Health Card Number:	
Emergency Contact Phone Number: ( ) -		Patient has private insurance coverage? <input type="checkbox"/> Yes <input type="checkbox"/> No	
PATIENT CLINICAL DETAILS			
Diagnosis:		Hemoglobin: _____ (g/L)	
Patient Weight (kg):		Ferritin: _____ (ng/mL)	
Date of Weight: (dd/mm/yyyy):		First Iron Infusion? <input type="checkbox"/> Yes <input type="checkbox"/> No	
Relevant Medical History/Notes:			
PRESCRIBER INFORMATION			
Prescribing Physician Name:		Prescribing Physician License #:	
Prescriber Phone Number: ( ) -		Address:	
Prescriber Fax Number: ( ) -		Email:	
PRESCRIPTION INFORMATION			
MONOFERRIC: Adult or Off label Pediatric *see pediatric section below			
<input type="checkbox"/> <b>Adult Protocol Infusion Dose</b> (select one): <input type="checkbox"/> 500 mg <input type="checkbox"/> 1000 mg <input type="checkbox"/> 1500 mg <input type="checkbox"/> Other: 20 mg/kg = _____ mg		<input type="checkbox"/> <b>Adult Protocol IV Bolus Injection</b> Administer _____ mg (up to 500 mg) of Monoferric once per week at a rate of up to 250 mg/minute.	
<b>Frequency</b> For each total dose (select one): <input type="checkbox"/> Administer as a single dose (if the total exceeds 20mg/kg or 1500mg, the dose must be split into two infusions at least one week apart. It is recommended, when possible, to give 20mg/kg in the first infusion) <input type="checkbox"/> Split dosing required (please specify): Day 0: _____mg, Day ____: _____mg Frequency of dose (select one): <input type="checkbox"/> One-time dose <input type="checkbox"/> Repeat Dosing Schedule: Administer # _____ infusions at a frequency of: <input type="checkbox"/> Weekly <input type="checkbox"/> Every 2 weeks <input type="checkbox"/> Monthly <input type="checkbox"/> Other: ____ Note: Prescribing physician is responsible for monitoring bloodwork and notifying Coverdale when no further treatment is required.			
<b>Off label:</b> <input type="checkbox"/> <b>Pediatric Protocol (SickKids Protocol below): Ordered Dose: _____ mg</b> If nausea or reaction occurs, nurse may administer: <b>DimenHYDRINATE 1 mg/kg (up to 50 mg) × 1 dose while in clinic</b> <b>Infuse slowly over 2 hours, titrating as tolerated.</b> Monitor vitals every 15 minutes. Post-infusion observation: 30 minutes (or per nursing judgement).			
<b>Frequency</b> For each total dose (select one): <input type="checkbox"/> Administer as a single dose (if the total exceeds 20mg/kg or 1500mg, the dose must be split into two infusions at least one week apart. It is recommended, when possible, to give 20mg/kg in the first infusion) <input type="checkbox"/> Split dosing required (please specify): Day 0: _____mg, Day ____: _____mg Frequency of dose (select one): <input type="checkbox"/> One-time dose <input type="checkbox"/> Repeat Dosing Schedule: Administer # _____ infusions at a frequency of: <input type="checkbox"/> Weekly <input type="checkbox"/> Every 2 weeks <input type="checkbox"/> Monthly <input type="checkbox"/> Other: ____ Note: Prescribing physician is responsible for monitoring bloodwork and notifying Coverdale when no further treatment is required.			
PRE-MEDICATIONS			
<input type="checkbox"/> None required <input type="checkbox"/> Other:			
FERINJECT: Adult or Pediatric			
<input type="checkbox"/> <b>Intravenous Infusion</b> (diluted in sterile 0.9% Sodium Chloride as per manufacturer's recommendations) <input type="checkbox"/> <b>Intravenous Injection</b> (undiluted) <b>Dose (Select one or write dose below):</b>			
<b>Determination of the Total Iron Need – Adults 18+</b>			
<b>Hb (g/L)</b>	<b>Bodyweight &lt;35 kg</b>	<b>Bodyweight 35 to &lt;70 kg</b>	<b>Bodyweight ≥70 kg</b>
<100	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 1500 mg	<input type="checkbox"/> 2000 mg
100 to <140	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 1000 mg	<input type="checkbox"/> 1500 mg
≥140	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 500 mg
<b>Determination of the Total Iron Need – Pediatric Ages 1 to 17 Years</b>			

# Iron (Monoferric, Ferinject and Venofer) Order Form \*TWO PAGES\*

Send completed form to Coverdale Infusion Clinics: Fax:1-888-236-3502 or by email: [enrollment@coverdaleclinic.com](mailto:enrollment@coverdaleclinic.com)

An infusion fee will apply and is payable by credit card to the Coverdale Resource Center at the time of booking. Please note that a cancellation fee may also apply.

Hb (g/L)	Bodyweight 10 kg	Bodyweight 20kg	Bodyweight 30kg	Bodyweight 40kg	Bodyweight 50kg	Bodyweight 60kg
70	<input type="checkbox"/> 300 mg	<input type="checkbox"/> 600 mg	<input type="checkbox"/> 900 mg	<input type="checkbox"/> 1200 mg	<input type="checkbox"/> 1350 mg	<input type="checkbox"/> 1500 mg
90	<input type="checkbox"/> 250 mg	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 750 mg	<input type="checkbox"/> 1100 mg	<input type="checkbox"/> 1200 mg	<input type="checkbox"/> 1350 mg
110	<input type="checkbox"/> 200 mg	<input type="checkbox"/> 400 mg	<input type="checkbox"/> 600 mg	<input type="checkbox"/> 900 mg	<input type="checkbox"/> 1000 mg	<input type="checkbox"/> 1100 mg
130	<input type="checkbox"/> 150 mg	<input type="checkbox"/> 300 mg	<input type="checkbox"/> 450 mg	<input type="checkbox"/> 700 mg	<input type="checkbox"/> 750 mg	<input type="checkbox"/> 800 mg
≥ 150	<input type="checkbox"/> 150 mg	<input type="checkbox"/> 300 mg	<input type="checkbox"/> 450 mg	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 500 mg	<input type="checkbox"/> 500 mg

☐ **Other:** \_\_\_\_\_ mg

**Based on the total iron requirement, administer FERINJECT as follows: Max single dose:** 15 mg iron/kg body weight or 1000 mg (adults)/750 mg (pediatric). **Max weekly dose:** 1000 mg (adults) and 750 mg (pediatric). **Pregnancy dosing (≥16 weeks):** **Max dose** – 1000 mg (Hb > 90), 1500 mg (Hb ≤ 90).

## Frequency:

☐ **One-time dose** (if dose is >1000mg (adults) or 750mg (pediatric), the dose must be split into two doses at least 7 days apart).

☐ **Split dosing required** (please specify): Day 0: \_\_\_\_\_ mg, Day \_\_\_\_: \_\_\_\_\_mg, Day \_\_\_\_: \_\_\_\_\_mg

Reassess Hb levels no sooner than 4 weeks after the final dose. If additional iron is needed, recalculate the required dose and submit a new medical order.

## PRE-MEDICATIONS

☐ None required ☐ Other:

## VENOFER: Adults or Off label Pediatric \*see pediatric section below

**Length of infusion:** Infuse IV over \_\_\_\_\_ minutes **Dose and route:** \_\_\_\_\_mg IV

**Frequency of dosing:** Administer every \_\_\_\_\_ (days/weeks/months) x \_\_\_\_\_ doses

## PRE-MEDICATIONS

☐ None required ☐ Other:

## Pregnancy Considerations:

**Is patient pregnant?** ☐ No ☐ Yes – If yes, by proceeding, you acknowledge that Monoferric, Venofer and Ferinject should only be used during pregnancy from gestation week 16 onward (specified in the Monoferric and Ferinject Product Monographs), when the benefit outweighs the risk to both mother and fetus, as per the Product Monograph. You have discussed the risks with the patient/guardian and wish to proceed.

**Pregnancy dosing (≥16 weeks):** **Max dose** – 1000 mg (Hb > 90), 1500 mg (Hb ≤ 90).

**Please specify infusion parameters for pregnant patients:**

**Start infusion slow for** \_\_\_\_\_ minutes, **slowly titrate up to infuse over** \_\_\_\_\_ minutes **as tolerated.**

**Monitor vital signs every** \_\_\_\_\_ minutes. **Post-infusion monitoring:** \_\_\_\_\_ minutes.

## Pediatric Considerations:

**Pediatric Use Acknowledgement (MONOFERRIC and VENOFER):** As per the Monoferric/Venofer Product Monograph, this medication has not been studied in pediatric populations and is not authorized by Health Canada for pediatric use. By signing below, you confirm that the risks have been discussed with the patient/guardian and that you wish to proceed with treatment as ordered. **Please indicate any additional protocols:**

*I authorize **Coverdale Resource Centre** to act as my designated agent to forward this prescription to the pharmacy chosen by the patient named above. This prescription is the original order and is intended solely for the patient's selected pharmacy.*

**Prescriber Signature:**

**Date (dd/mm/yyyy):**